




1/14




A



B



C



D

FIG. 1






FIG. 2

DNA Sample Prep

adaptor ligation



PURIFY 20KB FRAGMENTS

FIG. 3

DNA Sample Prep

strand isolation




FIG. 4

DNA Sample Prep

circularization




FIG. 5

Single Molecule Configuration

DNA-protein complexes




FIG. 6

Single-Molecule Isolation

bind to coverslip





FIG. 7

Sequence hundreds of individual single molecules in each well of a microtiter plate



TATGAAAAATTTCCGGTTTAAGCGGTTTCGGTTCTTCGTCATACTTAATGTTTTTATTTAAATAACCCCTCTGAAAAAGGAAA
CGACAGGTGCTGAAAGCGAGGCTTTTGGCCCTCTGTCGTTTCCTTTCTCTGTTTTTGTCCGTGGAAAGAAATGGAAGTCAACAAAA
GCAGCTGGTGACATTTTCGGTCCGAGTATCCGTACCATTCAGAACTGCGAGGAACAGGGAAATGCCCGTTCTCGAGGCGGTGGCAAGG
GTAATGAGGTGCTTTATGACTCTGCCGCCGTCATAAATGGTATGCCGAAGGGATGCTGAAATTGAGAACGAAAAAGCTGCCCGGGAG

10² PNACs per well

FIG. 8




FIG. 9A

10/14





FIG. 9B

11/14




FIG. 9C




FIG. 10




FIG. 11




FIG. 12